

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

**EP 1 357 112 A1**

(12)

**EUROPEAN PATENT APPLICATION**  
published in accordance with Art. 158(3) EPC

(43) Date of publication:  
29.10.2003 Bulletin 2003/44

(51) Int Cl.7: **C07D 221/14, C07D 311/08,  
C07D 311/12, C07D 311/80**

(21) Application number: **01970133.3**

(86) International application number:  
**PCT/JP01/08120**

(22) Date of filing: **19.09.2001**

(87) International publication number:  
**WO 02/051797 (04.07.2002 Gazette 2002/27)**

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE TR**

(30) Priority: **26.12.2000 JP 2000394669**


(71) Applicants:  
• **Applied Biosystems Japan Ltd.**  
**Tokyo 104-0032 (JP)**  
• **Ikeda, Hisafumi**  
**Nagareyama-shi, Chiba 270-0115 (JP)**

(72) Inventors:  
• **IKEDA, Hisafumi, Esuteeto Edogawadai 8-307**  
**Nagareyama-shi, Chiba 270-0115 (JP)**  
• **SAITO, Isao**  
**Kyoto-shi, Kyoto 607-8242 (JP)**  
• **KITAGAWA, Fumihiko**  
**Nagoya-shi, Aichi 452-0802 (JP)**


(74) Representative: **von Föner, Nicolai, Dr. et al**  
**v. Föner Ebbinghaus Finck Hano**  
**Patentanwälte**  
**Mariahilfplatz 2&3**  
**81541 München (DE)**

(54) **NOVEL FUNCTIONAL PEPTIDE NUCLEIC ACID MONOMER AND PROCESS FOR  
PRODUCING THE SAME**


(57) A compound represented by general formula (I) below;



(in the formula, A denotes




B denotes




R denotes H,  $\text{NO}_2$ ,  $\text{NH}_2$ ,  $\text{NHCbz}$ , Br, F, Cl or  $\text{SO}_3\text{Na}_2$ , and n is an integer of 1 to 4), and a process for producing the above compound characterised in that it includes a reaction between an activated ester and a *t*-butoxycarbonylami-noethylamine or an  $\omega$ -amino acid derivative.





Michael Egholm, Ole Buchardt, Peter E. Nielsen, and Rolf H. Berg  
*J. Am. Chem. Soc.* 1992, 114, 1895-1897.

after which a more efficient synthetic method:





Kim L. Duholm, Michael Egholm, et. al,  
*J. Org. Chem.* 1994, 59, 5767-5773.

was established.

[0009] However, as described above, since the Fmoc type, which is easy to handle, has been developed, the use of the tBoc type is becoming less frequent.

[0010] However, when introducing a functional molecule other than the four nucleic acids guanine, thymine, cytosine and adenine, for example, when introducing a photofunctional molecule, since the functional molecule that is to be introduced is often unstable under alkaline conditions, the tBoc type PNA backbone structure, which does not employ alkaline conditions, is very useful. With regard to a "process for producing t-butoxycarbonylaminoethylamines and amino acid derivatives", there is already a patent application filed by the present inventors as Japanese Patent Application No. 2000-268638.

[0011] Other than the above process, 5 examples of the synthesis of a monomer unit for a photofunctional PNA oligomer are known. All these cases employ the above-mentioned route, but their yields are either not described or very low (Peter E. Nielsen, Gerald Haaiman, Anne B. Eldrup PCT Int. Appl. (1998) WO 985295 A1 19981126, T. A. Tran, R.-H. Mattem, B. A. Morgan (1999) J. Pept. Res, 53, 134-145, Jesper Lohse et al. (1997) Bioconjugate Chem., 8, 503-509, Hans-georg Batz, Henrik Frydenlund Hansen, et al. PCT Int. Appl. (1998) WO 9837232 A2 19980827, Bruce Armitage, Troels Koch, et al. (1998) Nucleic Acid Res., 26, 715-720, Hans-georg Batz, Henrik Frydenlund Hansen, et al.). Furthermore, since the structures of the compounds used have the characteristic of being comparatively stable under alkaline conditions, it is surmised that, when a chromophore that is unstable under alkaline conditions is present, an efficient synthesis by a method similar to the above-mentioned conventional method, that is to say, route A below, is not possible.




[0012] There is therefore a strong desire for the establishment of a technique that functionalises a PNA monomer efficiently as well as for the development of a functional PNA monomer such as, for example, a photofunctional PNA

monomer.


**Disclosure of Invention**

[0013] It is an object of the present invention to provide a novel functional PNA monomer and an efficient synthetic method therefor that can eliminate the above-mentioned problems.

[0014] As a result of an intensive investigation by the present inventors in order to solve the above-mentioned problems, as shown in route B below, a photofunctional PNA monomer 4 has been synthesised successfully and substantially stoichiometrically using a *t*-butoxycarbonylaminoethylamine derivative 6 as a PNA backbone structure by condensation with an activated ester derivative 5 containing a pentafluorophenyl group 1.




[0015] Furthermore, the present inventors have also succeeded in synthesising a photofunctional PNA monomer 4 substantially stoichiometrically using an  $\omega$ -amino acid derivative 8 as a PNA backbone structure by condensation with an activated ester derivative 7 containing a pentafluorophenyl group 1 as shown in route C below:




[0016] The present inventors have found that the above-mentioned problems can be solved by the above-mentioned routes B and C, and the present invention has thus been accomplished.


[0017] That is to say, the present invention relates to a compound represented by general formula (I) below.




(In the formula, A denotes




B denotes



R denotes H, NO<sub>2</sub>, NH<sub>2</sub>, NHCbz, Br, F, Cl or SO<sub>3</sub>Na<sub>2</sub>, and n is an integer of 1 to 4, wherein when A denotes



then B denotes



).

[0018] Furthermore, the present invention relates to a process for producing a functional PNA monomer by reacting a *t*-butoxycarbonylaminoethylamine with a derivative of a functional molecule so as to incorporate the functional molecule into a PNA monomer, characterised in that the derivative of a functional molecule is an activated ester.

[0019] Furthermore, the present invention relates to the above-mentioned production process, characterised in that the activated ester has, on the carbonyl carbon forming the ester bond, a group represented by general formula (II) below:




40 (In the formula, A denotes

45

50

55





R denotes H, NO<sub>2</sub>, NH<sub>2</sub>, NHCbz, Br, F, Cl or SO<sub>3</sub>Na<sub>2</sub>, and n is an integer of 1 to 4.)

[0020] Furthermore, the present invention relates to the above-mentioned production process, characterised in that the activated ester has, on its carbonyl carbon, a pentafluorophenoxy group or a succinimidoxy group.

[0021] Furthermore, the present invention relates to a process for producing an activated ester, characterised in that it includes reacting a carboxylic acid derivative of a functional molecule with a compound having a pentafluorophenoxy group or a succinimidoxy group.


[0022] Furthermore, the present invention relates to a process for producing a carboxylic acid derivative of a functional molecule, characterised in that it includes reacting a derivative of the functional molecule with an aliphatic carboxylic acid.

[0023] Furthermore, the present invention relates to a process for producing a functional PNA monomer from a functional molecule, including:

- producing a derivative of the functional molecule from the functional molecule,
- producing a carboxylic acid derivative of the functional molecule from the derivative of the functional molecule,
- producing an activated ester from the carboxylic acid derivative of the functional molecule, and
- producing a functional PNA monomer from the activated ester, characterised in that it includes one or more of a) to c) below:

- a) in the production of the carboxylic acid derivative of the functional molecule, reacting the derivative of the functional molecule with an aliphatic carboxylic acid;
- b) in the production of the activated ester, reacting the carboxylic acid derivative of the functional molecule with a compound having a pentafluorophenoxy group or a succinimidoxy group; and
- c) in the production of the functional PNA monomer, reacting a t-butoxycarbonylaminoethylamine with the activated ester derivative of the functional molecule.

**[0024]** Furthermore, the present invention relates to a process for producing a functional PNA monomer by reacting a derivative of a functional molecule with an  $\omega$ -amino acid derivative represented by general formula (III) below, thereby incorporating the functional molecule into a PNA monomer, characterised in that the derivative of the functional molecule is an activated ester:




(In the formula,  $\text{R}^1$  denotes a hydrogen atom or a straight- or branched-chain  $\text{C}_1$  to  $\text{C}_5$  alkyl group, and  $m$  denotes an integer of 1 to 11.).

**[0025]** Furthermore, the present invention relates to the above-mentioned production process, characterised in that the activated ester has, on its carbonyl group forming an ester bond, a group represented by general formula (II) below, either directly or via an aliphatic chain or a peptide chain:



(In the formula, A denotes



R denotes H, NO<sub>2</sub>, NH<sub>2</sub>, NHCbz, Br, F, Cl or SO<sub>3</sub>Na<sub>2</sub>, and n is an integer of 1 to 4.).

[0026] Furthermore, the present invention relates to the above-mentioned production process, characterised in that the activated ester has, on its carbonyl carbon, a pentafluorophenoxy group or a succinimidoxy group.


[0027] Furthermore, the present invention relates to a process for producing a functional PNA monomer from a functional molecule, including producing an activated ester from the functional molecule and producing a functional PNA monomer from the activated ester, characterised in that

the production of the activated ester from the functional molecule includes reacting *m*-methyl red with a compound that contains a succinimidoxy group, and/or

the production of the functional PNA monomer from the activated ester includes reacting a benzyloxycarbonyl- $\omega$ -amino acid derivative represented by general formula (III) with the activated ester derivative of the functional molecule, thereby incorporating the functional molecule into a PNA monomer.


[0028] Features of the present invention are now explained in detail by comparing the processes of the present invention with the conventional processes.

[0029] The synthesis of tBoc type PNA monomer unit 4 usually employs route A below:



[0030] That is to say, it is a process in which, after synthesising 3 by dehydration condensation between 1 and 2, 3 is subjected to alkaline hydrolysis to give 4. This process is also used when introducing the four nucleic acid base groups guanine, thymine, cytosine, and adenine. When introducing a functional molecule (this refers to a known compound cited in Table 1) other than the above-mentioned groups, route A is also employed. However, since many photofunctional molecules are generally unstable under alkaline conditions, it is impossible to obtain 4 efficiently by route A. Instead of using the conventional PNA backbone structure 2, the condensation is therefore carried out using 6 obtained by the prior hydrolysis of 2. Since 6 includes a free carboxylic acid group, as is the case for 1, there is a possibility that an intramolecular dehydration condensation might occur, and a direct dehydration condensation using a condensing agent such as DCC cannot therefore be employed. The reaction is therefore devised so that the free carboxylic acid group and secondary amino group of 6 do not condense intramolecularly with each other by converting 1 into an activated ester derivative 5 using pentafluorophenol and then reacting 5 with 6 (route B). This gives 4 stoichiometrically. As hereinbefore described, there is no precedent for the process of synthesising 4 by converting a photofunctional molecule into its activated ester and then reacting it with 6, and it can be said that this process will hereafter be an indispensable technique for the synthesis of a variety of photofunctional PNA monomers.

[0031] Since benzyloxycarbonyl- $\omega$ -amino acid derivatives represented by general formula (III) below, which are used in route C, already have a linker (carboxylamino acid):



(in the formula,  $R^1$  denotes a hydrogen atom or a straight- or branched-chain  $C_1$  to  $C_5$  alkyl group, and  $m$  denotes an integer of 1 to 11),

the derivatives are very versatile, and reaction of the derivatives with an activated ester derivative gives a target functional PNA monomer unit in one step. Moreover, many commercially available benzyloxycarbonyl- $\omega$ -amino acid derivatives can be used. The route C is therefore particularly effective when targeting a comparatively expensive photofunctional molecule.

[0032] On the other hand, incorporation of a photofunctional molecule such as a sulphonyl chloride type or the highly sterically hindered methyl red into a PNA monomer can be carried out desirably by route B.

[0033] It is therefore possible to synthesise a variety of functional PNA monomer units by appropriately choosing one from the synthetic methods via route B and that via route C of the present invention.

[0034] In accordance with the present invention, photofunctional monomer units such as a Naphthalimide type, a Flavin type, a Dabcyl type, a Biotin type, an FAM type, a Rhodamine type, a TAMRA type, an ROX type, an HABA type, a Pyrene type and a Coumarin type can be obtained. It is also possible to obtain photofunctional monomer units other than the above-mentioned types and to obtain functional monomer units other than the photofunctional monomer units.

### Brief Description of Drawings


[0035]

FIG. 1 is a diagram showing differences in the structure and the electrical charge between DNA and PNA.

FIG. 2 is a diagram showing the structures of two types of PNA monomer unit.


### Modes for Carrying Out the Invention

[0036] Conventionally, in the synthesis of a photofunctional monomer unit, for example, in the synthesis of compound 4a,



a carboxylic acid derivative 1a of a functional molecule and a PNA backbone structure 2a are subjected to dehydration condensation to give 3a, which is then subjected to alkaline hydrolysis to give the target 4a (route A). Although the flavin skeleton of 1a is stable towards acid, it is easily decomposed under alkaline conditions to give 6,7-dimethylquinoxalinedione, and it is therefore impossible to obtain 4a efficiently even when 3a can be synthesised. When 1a is converted into an activated ester derivative 5a and then reacted with 6a, the reaction proceeds substantially stoichiometrically to give 4a with a yield of 85% (route B).

[0037] In the synthesis of compound 4b,



although a naphthalimide derivative 3b is obtained from the corresponding 1b and 2b with a yield of 44% the subsequent alkaline hydrolysis gives 4b in only 4% (route A). When 1b is converted into activated ester derivative 5b and then reacted with 6b, the reaction proceeds to give 4b with a yield of 75% (route B).

[0038] The production of 1a and 1b, which are carboxylic acid derivatives, is carried out using an aliphatic carboxylic acid, and preferably a straight-chain aliphatic carboxylic acid.

[0039] In the production of a monomer that is incorporated at a terminal amino group of a PNA, an activated ester having a succinimide group can preferably be used.

[0040] As the  $\omega$ -amino acid derivative represented by general formula (III) that is used in route C, one having a  $C_1$  to  $C_{11}$  carbon chain on the carbonyl carbon of its amino acid moiety can be used, but since a PNA is generally expected to be a hybrid with DNA, it is desirable for the derivative to be similar sterically to DNA. Taking this point into consideration, when a carboxylamino acid is used as a linker, one having Z-glycine as an amino moiety is the most suitable.

[0041] With regard to the method for synthesising a photofunctional PNA oligomer, the Fmoc method and the tBoc method can be used. The Fmoc method involves a protecting group removal step using an alkaline reagent, and its use is inappropriate when designing a photofunctional PNA oligomer. On the other hand, since the tBoc method does not employ alkaline conditions in its synthetic steps, it is suitable as a synthetic method for a photofunctional PNA oligomer. Application of the PNA monomer related to the present invention to the tBoc method therefore allows a photofunctional PNA oligomer to be synthesised efficiently.

## Examples

[0042] The present invention is explained below in more detail by way of examples, but the present invention is in no way limited thereby.

### (Example 1)

Synthesis of 2,3,4,5,6-pentafluorophenyl 2-(5,7,8-trimethyl-1,3-dioxo-2,5-dihydro-2,4-diazaphenazin-2-yl)acetate (5a)

[0043] EDC (73.2 mg, 382  $\mu$ mol) was added to a DMF solution (10 ml) of 1a (100 mg, 318  $\mu$ mol) and PfpOH (70.2 mg, 381  $\mu$ mol) at 0°C, and this reaction mixture was stirred at 0°C for 1 hour and at room temperature for 12 hours. This reaction mixture was concentrated under reduced pressure, and the residue was subjected to partition extraction with a water-chloroform system. The organic layer was dried with magnesium sulphate and concentrated under reduced pressure, and the residue was then purified by silica gel column chromatography (2.5% MeOH/ $CHCl_3$ ) to give 5a (130

mg, 85%).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  3.07 (s, 1 H), 7.44 (s, 1 H), 5.21 (s, 2 H), 4.14 (s, 3 H), 2.45 (s, 3 H); HRMS (FAB $^+$ , NBA/ $\text{CH}_2\text{Cl}_2$ )  $\text{C}_{21}\text{H}_{19}\text{N}_4\text{F}_5$  [(M+H) $^+$ ] Calcd. 481.0934, Exp. 481.0950;  $\text{UV } \lambda_{\text{max}}$  (DMF) 390, 460 (nm).

(Example 2)

Synthesis of 2-(N-(2-((t-butoxy)carbonylamino)ethyl))-2-(5,7,8-trimethyl-1,3-dioxo(2,5-dihydro-2,4-diazaphenazin-2-yl)acetyl)amino)acetic acid (4a)

**[0044]** Diisopropylethylamine (36.3  $\mu\text{L}$ , 208  $\mu\text{mol}$ ) was added to a DMF solution (10 mL) of 5a (100 mg, 208  $\mu\text{mol}$ ) and 6 (45.4 mg, 208  $\mu\text{mol}$ ) and stirred at room temperature for 15 hours. This was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (10-50% MeOH/ $\text{CHCl}_3$ ) to give 4a (130 mg, 85%).  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ )  $\delta$  7.94 (s) and 7.86 (s) (1 H), 7.80 (s) and 7.75 (s) (1 H), 5.03 (s) and 4.88 (s) (2 H), 4.17 (s) and 4.13 (s) (3 H), 3.64 and 3.52 (2 H), 3.38 and 3.26 (2 H), 2.58 (s) and 2.56 (s) (3 H), 2.46 (s) and 2.44 (s) (3 H), 1.46 (s) and 1.41 (s) (9 H); HRMS (FAB $^+$ , NBA/ $\text{CH}_2\text{Cl}_2$ )  $\text{C}_{24}\text{H}_{31}\text{O}_7\text{N}_6$  [(M+H) $^+$ ] Calcd. 515.2252, Exp. 515.2273;  $\text{UV } \lambda_{\text{max}}$  (DMF) 390, 460 (nm).

(Example 3)

Synthesis of N-(4-dimethylaminoazobenzene-2'-carbonyl)glycine (1c)

**[0045]** Triethylamine (732  $\mu\text{L}$ , 5.25 mmol) was added to a DMF solution (10 mL) of methyl red (1.35 g, 5 mmol) and t-butylglycine hydrochloride (880 mg, 5.25 mmol), while ice cooling DCC (1.13g, 5.5 mmol) was added thereto and this was stirred for 30 minutes and further at room temperature for 15 hours. The reaction mixture was filtered, the filtrate was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (0-10% acetone/ $\text{CH}_2\text{Cl}_2$ ) to give the t-butyl ester derivative (1.05 g, 55%) of 1c as orange needle-shaped crystals. This derivative (765 mg, 2 mmol) was added to formic acid (50 mL), stirred at room temperature for 2 days, and concentrated under reduced pressure to remove the formic acid, and the residue was purified by silica gel column chromatography (0-5% acetone/ $\text{CH}_2\text{Cl}_2$ ) to give 1c (549 mg, 84%) as red needle-shaped crystals.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  9.99 (brt, 1 H), 8.40 (d, J = 8 Hz, 1 H), 7.89 (d, J = 9 Hz, 2 H), 7.84 (d, J = 8 Hz, 1 H), 7.56 (t, J = 8 Hz, 1 H), 7.49 (t, J = 8 Hz, 1 H), 6.75 (d, J = 9 Hz, 2 H), 4.42 (d, J = 5 Hz, 2 H), 3.10 (s, 6 H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  167.61, 153.39, 150.86, 143.30, 132.49, 131.47, 129.38, 127.69, 126.28, 116.24, 111.63, 43.20, 40.24.

(Example 4)

Synthesis of 2,3,4,5,6-pentafluorophenyl N-(4-dimethylaminoazobenzene-2'-carbonyl)glycinate (5c)


**[0046]** DCC (308 mg, 1.5 mmol) was added to a DMF solution (10 mL) of 1c (326 mg, 1 mmol) and PfpOH (276 mg, 1.5 mmol) while ice cooling, and this reaction mixture was stirred at room temperature for 15 hours. The reaction mixture was filtered, the filtrate was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (0-5% acetone/ $\text{CH}_2\text{Cl}_2$ ) to give 5c (449 mg, 91%) as an orange powder.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  10.14 (brt, 1 H), 8.37 (d, J = 8 Hz, 1 H), 7.78 (d, J = 9 Hz, 2 H), 7.76 (d, J = 8 Hz, 1 H), 7.57 (t, J = 8 Hz, 1 H), 7.50 (t, J = 8 Hz, 1 H), 6.74 (d, J = 9 Hz, 2 H), 4.68 (d, J = 5 Hz, 2 H), 3.06 (s, 6 H).

(Example 5)

Synthesis of 2-(N-(2-((t-butoxy)carbonylamino)ethyl))-2-(4-dimethylaminoazobenzene-2'-carbonylamino)acetyl)amino)acetic acid (4c)

**[0047]** Diisopropylethylamine (85  $\mu\text{L}$ , 0.5 mmol) was added to a DMF solution (5 mL) of 5c (246 mg, 0.5 mmol) and 6 (109 mg, 0.5 mmol) and stirred at room temperature for 15 hours. This was concentrated under reduced pressure and the residue was purified by silica gel column chromatography (0-30% MeOH/ $\text{CH}_2\text{Cl}_2$ ) to give 4c (225 mg, 72%).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  9.99 (s) and 9.85 (s) (1 H), 8.3-7.6 (m, 4 H), 7.4-7.2 (m, 2 H), 6.67 (s) and 6.59 (s) (2 H), 5.62 (s) and 5.27 (s) (1 H), 4.35 (s) and 4.20 (s) (2 H), 3.99 and 3.90 (2 H), 3.5 (brs) and 3.3 (brs) (2 H), 3.2 (brs) and 3.0 (brs) (2 H), 2.99 (s) and 2.87 (s) (6 H), 1.25 (brs, 9 H).

**[0048]** The route from 1c to 4c was as follows:



(Example 6)

Synthesis of *N*-(4-hydroxyazobenzene-2'-carbonyl)glycine (1d)

50 [0049] Triethylamine (732  $\mu$ L, 5.25 mmol) was added to a DMF solution (10 mL) of HABA (1.21 g, 5 mmol) and *t*-butylglycine hydrochloride (880 mg, 5.25 mmol), while ice cooling DCC (1.13 g, 5.5 mmol) was then added and this was stirred for 30 minutes and further at room temperature for 15 hours. The reaction mixture was filtered, the filtrate was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (0-5% acetone/ $\text{CH}_2\text{Cl}_2$ ) to give the *t*-butyl ester derivative (1.73 g, 97%) of 1d as an orange powder. This (1.07 g, 3 mmol)

55 was added to formic acid (50 mL), stirred at room temperature for 2 days, and concentrated under reduced pressure to remove the formic acid, and the residue was purified by silica gel column chromatography (0-5% acetone/ $\text{CH}_2\text{Cl}_2$ ) to give 1d (0.89 g, 99%) as an orange powder.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  10.45 (brt, 1 H), 8.88 (brt,  $J$  = 5 Hz, 1 H), 7.91 (d,  $J$  = 9 Hz, 2 H), 7.85 (d,  $J$  = 8 Hz, 1 H), 7.70 (d,  $J$  = 8 Hz, 1 H), 7.60 (t,  $J$  = 8 Hz, 1 H), 7.57 (t,  $J$  = 8 Hz, 1-H), 6.94 (d,  $J$



= 9 Hz, 2 H), 4.05 (d, J = 7.7, 2 H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  171.18, 166.44, 161.54, 159.13, 145.34, 133.20, 131.09, 130.18, 129.61, 125.86, 115.88, 41.60.

(Example 7)

Synthesis of 2,3,4,5,6-pentafluorophenyl *N*-(4-hydroxyazobenzene-2'-carbonyl)glycinate (5d)






[0050] DCC (308 mg, 1.5 mmol) was added to a DMF solution (10 mL) of 1d (299 mg, 1 mmol) and PfpOH (276 mg, 1.5 mmol) while ice cooling, and this reaction mixture was stirred at room temperature for 15 hours. The reaction mixture was filtered, the filtrate was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (0-5% acetone/ $\text{CH}_2\text{Cl}_2$ ) to give 5d (46 mg, 10%) as an orange powder.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  9.67 (brs, 1 H), 9.02 (brt, 1 H), 7.8-7.7 (m, 3 H), 7.6-7.5 (m, 2 H), 7.23 (d, J = 9 Hz, 1H), 6.86 (d, J = 9 Hz, 2 H), 4.67 (d, J = 5 Hz, 2 H).

(Example 8)

Synthesis of 2-(*N*-(2-((*t*-butoxy)carbonylamino)ethyl)-2-(4-hydroxyazobenzene-2'-carbonylamino)acetyl amino) acetic acid (4d)

[0051] Diisopropylethylamine (14  $\mu\text{L}$ , 80  $\mu\text{mol}$ ) was added to a DMF solution (5 mL) of 5d (37 mg, 80  $\mu\text{mol}$ ) and 6 (18 mg, 80  $\mu\text{mol}$ ) and stirred at room temperature for 15 hours. This was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (0-30% MeOH/ $\text{CH}_2\text{Cl}_2$ ) to give 4d (13 mg, 33%).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  -9.77 (s) and 9.59 (s) (1 H), 8.26 (s) and 8.14 (s) (2H), 7.9-7.6 (m, 2 H), 7.6-7.3 (m, 2 H), 7.0-6.6 (m, 2 H), 5.35 (s) and 5.05(s) (1 H), 4.40 (s) and 4.24 (s) (2 H), 3.98 (s, 2 H), 3.6-3.3 (m, 2 H), 3.21 (s) and 3.02 (s) (2 H), 1.28 (s) and 1.18 (s) (9 H).

[0052] The route from 1d to 4d was as follows.



(Example 9)

#### Synthesis of FAM-Gly-BocPNA-OH

45

[0053] 5,6-FAM *N*-hydroxysuccinimide ester (50 mg, 0.11 mmol) and triethylamine (250  $\mu$ L, 2.0 mmol) were added in that order to a dimethylformamide solution (5 mL) of Gly-BocPNA-OH (30.3 mg, 0.10 mmol) and stirred at room temperature for 15 hours. After completion of the reaction, the mixture was concentrated under reduced pressure, and the residue was subjected to silica gel column chromatography (0-25% MeOH/dichloromethane) to give FAM-Gly-BocPNA-OH (69.8 mg, 100%) as a yellow powder. FABMS  $m/z$  634  $[(M+H)^+]$ ; HRMS (FAB $^+$ ) calcd. for  $C_{32}H_{32}O_{11}N_3$   $[(M+H)^+]$  634.1959, observed 634.2034.

(Example 10)

#### Synthesis of TAMRA-Gly-BocPNA-OH

55

[0054] 5,6-TAMRA *N*-hydroxysuccinimide ester (5 mg, 9.5  $\mu$ mol) and triethylamine (20  $\mu$ L, 140  $\mu$ mol) were added in that order to a dimethylformamide solution (5 mL) of Gly-BocPNA-OH (5.8 mg, 21  $\mu$ mol) and stirred at room temperature

for 15 hours. After completion of the reaction, the mixture was concentrated under reduced pressure, and the residue was subjected to silica gel column chromatography (0-30% MeOH/dichloromethane) to give TAMRA-Gly-BocPNA-OH (6 mg, 100%) as reddish purple powder. FABMS  $m/z$  688 [(M+H)<sup>+</sup>]; HRMS (FAB<sup>+</sup>) calcd. for C<sub>36</sub>H<sub>42</sub>O<sub>9</sub>N<sub>5</sub> [(M+H)<sup>+</sup>] 688.2904, observed 688.2993.

(Example 11)

#### Synthesis of ROX-Gly-BocPNA-OH

**[0055]** 5,6-ROX N-hydroxysuccinimide ester (5 mg, 8 μmol) and triethylamine (20 μL, 140 μmol) were added in that order to a dimethylformamide solution (5 mL) of Gly-BocPNA-OH (4.9 mg, 18 μmol) and stirred at room temperature for 15 hours. After completion of the reaction, the mixture was concentrated under reduced pressure, and the residue was subjected to silica gel column chromatography (0-30% MeOH/dichloromethane) to give ROX-Gly-BocPNA-OH (6 mg, 100%) as a purple powder. FABMS  $m/z$  792 [(M+R)<sup>+</sup>]; HRMS (FAB<sup>+</sup>) calcd. for C<sub>44</sub>H<sub>50</sub>O<sub>9</sub>N<sub>5</sub> [(M+H)<sup>+</sup>] 792.3530, observed 792.3615.

(Example 12)

#### Synthesis of 2,3,4,5,6-pentafluorophenyl *N*-(4-dimethylaminoazobenzene-2'-carbonyl)glycinate (*o*-MR-Gly-Opfp)

**[0056]** DCC (308 mg, 1.5 mmol) was added to a DMF solution (10 mL) of *o*-MR-Gly-OH (326 mg, 1 mmol) and PfpOH (276 mg, 1.5 mmol) while ice cooling, and the reaction mixture was stirred at room temperature for 15 hours. The reaction mixture was filtered, the filtrate was concentrated under reduced pressure and the residue was purified by silica gel column chromatography (0-5% acetone/CH<sub>2</sub>Cl<sub>2</sub>) to give *o*-MR-Gly-Opfp (449 mg, 91%) as an orange powder. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 10.14 (brt, 1 H), 8.37 (d, J = 8 Hz, 1 H), 7.78 (d, J = 9 Hz, 2 H), 7.76 (d, J = 8 Hz, 1 H), 7.57 (t, J = 8 Hz, 1 H), 7.50 (t, J = 8 Hz, 1 H), 6.74 (d, J = 9 Hz, 2 H), 4.68 (d, J = 5 Hz, 2 H), 3.06 (s, 6 H).

(Example 13)

#### Synthesis of 2-(*N*-(2-((*tert*-butoxy)carbonylamino)ethyl)-2-(4-dimethylaminoazobenzene-2'-carbonylamino)acetylamin)acetic acid (*o*-MR-Gly-BocPNA-OH)

**[0057]** Diisopropylethylamine (85 μL, 0.5 mmol) was added to a DMF solution (5 mL) of *o*-MR-Gly-Opfp (246 mg, 0.5 mmol) and BocPNA-OH (109 mg, 0.5 mmol) and stirred at room temperature for 15 hours. This was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (0-30% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to give *o*-MR-Gly-BocPNA-OH (225 mg, 72%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 9.99 (s) and 9.85 (s) (1 H), 8.3-7.6 (m) (4 H), 7.4-7.2 (m) (2 H), 6.67 (s) and 6.59 (s) (2 H), 5.62 (s) and 5.27 (s) (1 H), 4.35 (s) and 4.20 (s) (2 H), 3.99 and 3.90 (2 H), 3.5 (brs) and 3.3 (brs) (2 H), 3.2 (brs) and 3.0 (brs) (2 H), 2.99 (s) and 2.87 (s) (6 H), 1.25 (brs, 9 H).

(Example 14)

#### Synthesis of 2-(*N*-(2-((*tert*-butoxy)carbonylamino)ethyl)-2-(4-dimethylaminoazobenzene-4'-carbonylamino)acetylamin)acetic acid (Dabcyl-Gly-BocPNA-OH (*p*-MR-Gly-BocPNA-OH))

**[0058]** Dabcyl *N*-hydroxysuccinimide ester (145 mg, 0.40 mmol) and triethylamine (600 μL, 4.5 mmol) were added in that order to a dimethylformamide solution (10 mL) of Gly-BocPNA-OH (100 mg, 0.39 mmol) and stirred at room temperature for 15 hours. After completion of the reaction, the mixture was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (0-4% MeOH/dichloromethane) to give Dabcyl-Gly-BocPNA-OH (184 mg, 90%) as a reddish brown powder. <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 8.18 (d, J = 7 Hz, 2 H), 7.91 (d, J = 7 Hz, 2 H), 7.88 (d, J = 7 Hz, 2 H), 6.77 (d, J = 7 Hz, 2 H), 5.76 (s) and 5.30 (s) (2 H), 4.22 (brs) and 4.05 (brs) (2 H), 3.73 (brs) and 3.49 (brs) (2 H), 3.47 (brs) and 3.29 (brs) (2 H), 1.26 (s, 9 H); FABMS  $m/z$  527 (M+H)<sup>+</sup>].

(Example 15)

#### Synthesis of *N*-(4-hydroxyazobenzene-2'-carbonyl)glycine (HABA-Gly-OH)

**[0059]** Triethylamine (732 μL, 5.25 mmol) was added to a DMF solution (10 mL) of HABA (1.21 g, 5 mmol) and *t*-butylglycine hydrochloride (880 mg, 5.25 mmol), while ice cooling DCC (1.13 g, 5.5 mmol) was then added thereto and

the mixture was stirred for 15 minutes and further at room temperature for 15 hours. The reaction mixture was filtered, the filtrate was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (0-5% Acetone/CH<sub>2</sub>Cl<sub>2</sub>) to give the *t*-butyl ester derivative of HABA-Gly-OH (1.73 g, 97%) as an orange powder. This (1.07 g, 3 mmol) was added to formic acid (50 mL) and stirred at room temperature for 2 days, concentrated under reduced pressure to remove the formic acid, and the residue was purified by silica gel column chromatography (0-5% acetone/CH<sub>2</sub>Cl<sub>2</sub>) to give HABA-Gly-OH (0.89 g, 99%) as an orange powder. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 10.45 (brt, 1 H), 8.88 (brt, J = 5 Hz, 1 H), 7.91 (d, J = 9 Hz, 2 H), 7.85 (d, J = 8 Hz, 1 H), 7.70 (d, J = 8 Hz, 1 H), 7.60 (t, J = 8 Hz, 1 H), 7.57 (t, J = 8 Hz, 1 H), 6.94 (d, J = 9 Hz, 2 H), 4.05 (d, J = 5 Hz, 2 H); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 171.18, 166.44, 161.54, 149.13, 145.34, 133.20, 131.09, 130.18, 129.61, 125.86, 116.00, 115.88, 41.60.

(Example 16)

Synthesis of 2,3,4,5,6-pentafluorophenyl *N*-(4-hydroxyazobenzene-2'-carbonyl)glycinate (HABA-Gly-Opfp)

**[0060]** DCC (308 mg, 1.5 mmol) was added to a DMF solution (10 mL) of HABA-Gly-OH (299 mg, 1 mmol) and PfpOH (276 mg, 1.5 mmol) while ice cooling, and the reaction mixture was stirred at room temperature for 15 hours. The reaction mixture was filtered, the filtrate was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (0-5% acetone/CH<sub>2</sub>Cl<sub>2</sub>) to give HABA-Gly-OPfp (46 mg, 10%) as an orange powder. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 9.67 (brs, 1 H), 9.02 (brt, 1 H), 7.8-7.7 (m, 3 H), 7.6-7.5 (m, 2 H), 7.23 (d, J = 9 Hz, 1 H), 6.86 (d, J = 9 Hz, 2 H), 4.67 (d, J = 5 Hz, 2 H).

(Example 17)

Synthesis of 2-(*N*-(2-((*tert*-butoxy)carbonylamino)ethyl)-2-(4-hydroxyazobenzene-2'-carbonylamino)acetyl amino) acetic acid (HABA-Gly-BocPNA-OH)

**[0061]** Diisopropylethylamine (14 mL, 80 μmol) was added to a DMF solution (5 mL) of HABA-Gly-OPfp (37 mg, 80 μmol) and BocPNA-OH (18 mg, 80 μmol) and stirred at room temperature for 15 hours. This was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (0-30% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to give HABA-Gly-BocPNA-OH (13 mg, 33%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 9.77 (s) and 9.59 (s) (1 H), 8.26 (s) and 8.14 (s) (2 H), 7.9-7.6 (m, 2 H), 7.6-7.3 (m, 2 H), 7.0-6.6 (m, 2 H), 5.35 (s) and 5.05 (s) (1 H), 4.40 (s) and 4.24 (s) (2 H), 3.98 (s, 2 H), 3.6-3.3 (m, 2 H), 3.21 (s) and 3.02 (s) (2 H), 1.28 (s) and 1.18 (s) (9 H).

(Example 18)

Synthesis of 2,3,4,5,6-pentafluorophenyl 2-(5,7,8-trimethyl-1,3-dioxo-2,5-dihydro-2,4-diazaphenazin-2-yl)acetate (Flavin-Opfp)

**[0062]** EDC (73.2 mg, 382 μmol) was added to a DMF solution (10 mL) of Flavin (100 mg, 318 μmol) and PfpOH (70.2 mg, 381 μmol) at 0°C, and the reaction mixture was stirred at 0°C for 1 hour and at room temperature for 12 hours. The reaction mixture was concentrated under reduced pressure, and the residue was subjected to partition extraction with a water-chloroform system. The organic layer was dried with magnesium sulphate and concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (2.5% MeOH/CHCl<sub>3</sub>) to give Flavin-OPfp (130 mg, 85%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.07 (s, 1 H), 7.44 (s, 1 H), 5.21 (s, 2 H), 4.14 (s, 3 H), 2.55 (s, 3 H), 2.45 (s, 3 H); HRMS (FAB<sup>+</sup>, NBA/CH<sub>2</sub>Cl<sub>2</sub>) calcd. for C<sub>21</sub>H<sub>14</sub>O<sub>4</sub>N<sub>4</sub>F<sub>5</sub> [(M+H)<sup>+</sup>] 481.0934, observed 481.0950; UV λ<sub>max</sub> (DMF) 390, 460 (nm).

(Example 19)

Synthesis of 2-(*N*-(2-((*tert*-butoxy)carbonylamino)ethyl)-2-(5,7,8-trimethyl-1,3-dioxo(2,5-dihydro-2,4-di-azaphenazin-2-yl)acetyl amino) acetic acid (Flavin-BocPNA-OH)

**[0063]** Diisopropylethylamine (36.3 μL, 208 μmol) was added to a DMF solution (10 mL) of Flavin-OPfp (100 mg, 208 μmol) and BocPNA-OH (45.4 mg, 208 μmol) and stirred at room temperature for 15 hours. This was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (10-50% MeOH/CHCl<sub>3</sub>) to give Flavin-BocPNA-OH (130 mg, 85%). <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 7.94 (s) and 7.86 (s) (1 H), 7.80 (s) and 7.75 (s) (1 H), 5.03 (s) and 4.88 (s) (2 H), 4.17 (s) and 4.13 (s) (3 H), 3.64 and 3.52 (2 H), 3.38 and 3.26 (2 H), 2.58 (s) and 2.56 (s) (3 H), 2.46 (s) and 2.44 (s) (3 H), 1.46 (s) and 1.41 (s) (9 H); HRMS (FAB<sup>+</sup>, NBA/CH<sub>2</sub>Cl<sub>2</sub>) calcd. for C<sub>24</sub>H<sub>31</sub>O<sub>7</sub>N<sub>6</sub>

[(M+H)<sup>+</sup>] 515.2252, observed 515.2273; UV  $\lambda_{\text{max}}$  (DMF) 390, 460 (nm).

(Example 20)

5 Synthesis of 2',3',4',5',6'-pentafluorophenyl 1,3-dioxo-1H-benz[de]isoquinoline-2(3H)-acetate (NI-Opfp)

[0064] DCC (155 mg, 0.75 mmol) was added to a DMF solution (5 mL) of NI-OH (192 mg, 0.75 mmol) and PfpOH (152 mg, 0.83 mmol) while ice cooling, and the reaction mixture was stirred at room temperature for 15 hours. The reaction mixture was filtered, the filtrate was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (CH<sub>2</sub>Cl<sub>2</sub>) to give NI-OPfp (277 mg, 87%) as a red powder. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  8.64 (d, J = 8 Hz, 2 H), 8.25 (d, J = 8 Hz, 2 H), 7.78 (t, J = 8 Hz, 2 H), 5.29 (s, 2 H).

(Example 21)

15 Synthesis of 2-(N-(2-((*tert*-butoxy)carbonylamino)ethyl)-2-(1,3-dioxo-1H-benz[de]isoquinoline-2(3H))acetylamino)acetic acid (NI-BocPNA-OH)

[0065] Diisopropylethylamine (87  $\mu$ L, 0.50 mmol) was added to a DMF solution (10 mL) of NI-OPfp (211 mg, 0.50 mmol) and BocPNA-OH (120 mg, 0.55 mmol) and stirred at room temperature for 15 hours. This was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (0-20% MeOH/CHCl<sub>3</sub>) to give NI-BocPNA-OH (130 mg, 85%). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  8.47 (m, 4 H), 7.86 (dd, J = 8.3, 7.3 Hz, 2 H), 4.73 (s, 2 H); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  167.84, 163.59, 134.20, 131.44, 131.39, 128.11, 126.78, 122.00, 61.59, 41.44, 14.29; HRMS (FAB<sup>+</sup>, NBA/CH<sub>2</sub>Cl<sub>2</sub>) calculated for (C<sub>16</sub>H<sub>13</sub>NO<sub>4</sub>)H<sup>+</sup> 284.2933, observed 456.1767; UV  $\lambda_{\text{max}}$  (DMF) 333 nm.

25 (Example 22)

Synthesis of 2',3',4',5',6'-pentafluorophenyl 1,3-dioxo-5-nitro-1H-benz[de]isoquinoline-2(3H)-acetate (NI (NO<sub>2</sub>)-Opfp)

[0066] EDC (73.2 mg, 382  $\mu$ mol) was added to a DMF solution (10 mL) of NI(NO<sub>2</sub>)-OH (100 mg, 318  $\mu$ mol) and PfpOH (70.2 mg, 381  $\mu$ mol) at 0°C, and the reaction mixture was stirred at 0°C for 1 hour and at room temperature for 12 hours. The reaction mixture was concentrated under reduced pressure, and the residue was subjected to partition extraction with a water-chloroform system. The organic layer was dried with magnesium sulphate and concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (2.5% MeOH/CHCl<sub>3</sub>) to give NI(NO<sub>2</sub>)-OPfp (130 mg, 85%). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  8.07 (s, 1 H), 7.44 (s, 1 H), 5.21 (s, 2 H), 4.14 (s, 3 H), 2.55 (s, 3 H), 2.45 (s, 3 H); HRMS (FAB<sup>+</sup>, NBA/CH<sub>2</sub>Cl<sub>2</sub>) calcd. for C<sub>21</sub>H<sub>14</sub>O<sub>4</sub>N<sub>4</sub>F<sub>5</sub> [(M+H)<sup>+</sup>] 481.0934, observed 481.0950; UV  $\lambda_{\text{max}}$  (DMF) 390, 460 (nm).

(Example 23)

40 Synthesis of 2-(N-(2-((*tert*-butoxy)carbonylamino)ethyl)-2-(1,3-dioxo-5-nitro-1H-benz[de]isoquinoline-2(3H))acetylamino)acetic acid (NI (NO<sub>2</sub>)-BocPNA-OH)

[0067] Diisopropylethylamine (36.3  $\mu$ L, 208  $\mu$ mol) was added to a DMF solution (10 mL) of NI(NO<sub>2</sub>)-OPfp (100 mg, 208  $\mu$ mol) and BocPNA-OH (45.4 mg, 208  $\mu$ mol) and stirred at room temperature for 15 hours. This was concentrated under reduced pressure and the residue was purified by silica gel column chromatography (10-50% MeOH/CHCl<sub>3</sub>) to give NI (NO<sub>2</sub>)-BocPNA-OH (130 mg, 85%). <sup>1</sup>H NMR (CD<sub>3</sub>OD)  $\delta$  7.94 (s) and 7.86 (s) (1 H), 7.80 (s) and 7.75 (s) (1 H), 5.03 (s) and 4.88 (s) (2 H), 4.17 (s) and 4.13 (s) (3 H), 3.64 and 3.52 (2 H), 3.38 and 3.26 (2 H), 2.58 (s) and 2.56 (s) (3 H), 2.46 (s) and 2.44 (s) (3 H), 1.46 (s) and 1.41 (s) (9 H); HRMS (FAB<sup>+</sup>, NBA/CH<sub>2</sub>Cl<sub>2</sub>) calcd. for C<sub>24</sub>H<sub>31</sub>O<sub>7</sub>N<sub>6</sub> [(M+H)<sup>+</sup>] 515.2252, observed 515.2273; UV  $\lambda_{\text{max}}$  (DMF) 390, 460 (nm).

50

(Example 24)

Synthesis of 5-acetylamino-1,3-dioxo-1H-benz[de]isoquinoline-2(3H)-acetic Acid (NI(NHAc)-OH)

55 [0068] NI (NH<sub>2</sub>)-OH (100 mg, 0.37 mmol) was dissolved in pyridine (3 mL) and AC<sub>2</sub>O (3 mL) and stirred at room temperature for 15 hours. The mixture was concentrated under reduced pressure, then washed with dichloromethane, filtered and dried to give NI(NHAc)-OH (103.2 mg, 89%). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  8.79 (s, 1 H), 8.61 (s, 1 H), 8.40 (d, J = 8 Hz, 1 H), 8.37 (d, J = 8 Hz, 1 H), 7.82 (t, J = 8 Hz, 1 H), 4.71 (s, 2 H), 2.16 (s, 3 H).

## (Example 25)

Synthesis of 2',3',4',5',6'-pentafluorophenyl 5-acetylamino-1,3-dioxo-1H-benz[de]isoquinoline-2(3H)-acetate (NI (NHAc)-OPfp)

[0069] DCC (71 mg, 0.34 mmol) was added to a DMF solution (5 mL) of NI(NHAc)-OH (97 mg, 0.31 mmol) and PfpOH (63 mg, 0.34 mmol) at 0°C and the reaction mixture was stirred at 0°C for 1 hour and at room temperature for 15 hours. The reaction mixture was filtered and then concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (0-20% acetone/CHCl<sub>3</sub>) to give NI (NHAc)-OPfp (140 mg, 95%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.96 (s, 1 H), 8.53 (d, J = 7.7 Hz, 1 H), 8.32 (d, J = 1.6 Hz, 1 H), 8.20 (d, J = 7.7 Hz, 1 H), (t, J = 7.5 Hz, 2 H), 7.82 (brs, 1 H), 7.74 (d, J = 7.7 Hz, 1 H), 5.27 (s, 2 H), 2.28 (s, 3 H).

## (Example 26)

Synthesis of 2-(N-(2-((*tert*-butoxy)carbonylamino)ethyl)-2-(5-acetylamino-1,3-dioxo-1H-benz[de]isoquinoline-2(3H)) acetylamino)acetic acid (NI (NHAc)-BocPNA-OH)

[0070] Diisopropylethylamine (54 μL, 0.30 mmol) was added to a DMF solution (5 mL) of NI (NHAc)-OPfp (140 mg, 0.29 mmol) and BocPNA-OH (67 mg, 0.30 mmol) and stirred at room temperature for 15 hours. This was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (2-20% MeOH/CHCl<sub>3</sub>) to give NI (NHAc) - BocPNA-OH (117 mg, 85%). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.4-7.3 (m, 5 H), 5.05 (brs) and 4.90 (brs) (2 H), 3.76 (brs) and 3.54 (brs) (2 H), 3.64 (s) and 3.49 (s) (2 H), 3.54 (brs) and 3.41 (brs) (2 H), 2.15 (s) and 2.04 (s) (3 H), 1.48 (s) and 1.45 (s) (9 H).

## (Example 27)

Synthesis of succinimidyl N-4-dimethylaminoazobenzene-3'-carbonate (*m*-MR-OSu)

[0071] DCC (100 mg, 0.50 mmol) was added to a DMF solution (7 mL) of *m*-methyl Red (*m*-MR-OH; 110 mg, 0.41 mmol) and N-hydroxysuccinimide (60 mg, 0.52 mmol) at 0°C, and the reaction mixture was stirred for 30 minutes and then at room temperature for 15 hours. The reaction mixture was filtered and distilled under reduced pressure, and the residue was subjected to silica gel column chromatography (CH<sub>2</sub>Cl<sub>2</sub>) to give *m*-MR-OSu (124 mg; 82%) as an orange powder. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.59 (s, 1 H), 8.13 (t, J = 9.1 Hz, 2 H), 7.90 (d, J = 9.1 Hz, 2 H), 7.61 (t, J = 7.9 Hz, 1 H), 6.77 (d, J = 9.1 Hz, 2 H), 3.11 (s, 6 H), 2.93 (brs, 4 H).

## (Example 28)

Synthesis of 2-(N-(2-((*tert*-butoxy)carbonylamino)ethyl)-2-(4-dimethylaminoazobenzene-3'-carbonylamino) acetylamino)acetic acid (*m*-MR-Gly- BocPNA-OH)

[0072] *m*-MR-OSu (73 mg, 0.20 mmol) and triethylamine (350 μL, 2.7 mmol) were added in that order to a DMF solution (10 mL) of Gly-BocPNA-OH (50 mg, 0.18 mmol) and stirred at room temperature for 15 hours. After completion of the reaction, the mixture was concentrated under reduced pressure, and the residue was subjected to silica gel column chromatography (0-10% MeOH/dichloromethane) to give *m*-MR-Gly-BocPNA-OH (95 mg, 100%) as an orange powder. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 8.26 (s, 1 H), 7.92 (d, J = 7.6 Hz, 2 H), 7.83 (d, J = 9.1 Hz, 2 H), 7.62 (t, J = 7.6 Hz, 1 H), 6.88 (brt) and 6.74 (brt) (1 H), 6.85 (d, J = 9.1 Hz, 2 H), 4.22 (d, J = 2.7 Hz, 2 H), 3.99 (s) and 3.89 (s) (2 H), 3.44 (t, J = 6.4 Hz, 1 H), 3.4-3.25 (brs, 4 H), 3.07 (s, 6 H), 1.39 (s) and 1.37 (s) (9 H).

## (Example 29)

Synthesis of 2-(N-(2-((*tert*-butoxy)carbonylamino)ethyl)-N'-((1-pyrenyl-*n*-butyl)glycyl))acetic acid (Pyrene-Gly-BocPNA-OH)

[0073] Pyrene-OSu (39 mg, 0.10 mmol) and triethylamine (138 μL, 1.0 mmol) were added in that order to a DMF solution (5 mL) of Gly-BocPNA-OH (25 mg, 0.09 mmol) and stirred at room temperature for 15 hours. After completion of the reaction, the mixture was concentrated under reduced pressure, and the residue was subjected to silica gel column chromatography (0-10% MeOH/dichloromethane) to give Pyrene-Gly-BocPNA-OH (30 mg, 61%) as a pale yellow powder. HRMS (FAB<sup>+</sup>) calcd. for C<sub>31</sub>H<sub>35</sub>O<sub>6</sub>N<sub>3</sub>Na [(M+Na)<sup>+</sup>] 568.2526, observed 568.2429.

(Example 30)

Synthesis of 2-(N-(2-((*tert*-butoxy)carbonylamino)ethyl)-2-(7-diethylaminocoumarin-3-carbonyl)glycyl)acetic acid (Coumarin-Gly-BocPNA-OH)


[0074] Coumarin-OSu (15 mg, 0.042 mmol) and triethylamine (55.5  $\mu$ L, 0.4 mmol) were added in that order to a DMF solution (5 mL) of Gly-BocPNA-OH (12.7 mg, 0.046 mmol) and stirred at room temperature for 15 hours. After completion of the reaction, the mixture was concentrated under reduced pressure, and the residue was subjected to silica gel column chromatography (0-20% MeOH/dichloromethane) to give Coumarin-Gly-BocPNA-OH (23 mg, 100%) as a yellow powder.  $^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  8.68 (s) and 8.66 (s) (1 H), 7.70 and 7.69 (each d,  $J = 9.1$  Hz) (1 H), 6.89 (brt) and 6.75 (brt) (1 H), 6.80 (d,  $J = 9.1$  Hz, 1 H), 6.62 (s, 1 H), 4.25 (brd) and 4.07 (brd) (2 H), 4.13 (m, 1 H), 3.98 (s) and 3.89 (s) (2 H), 3.48 (q,  $J = 6.8$  Hz, 4 H), 3.35 (m, 2 H), 3.13 (brq) and 3.07 (brq) (2 H), 1.37 (s) and 1.36 (s) (9 H), 1.14 (t,  $J = 6.8$  Hz, 6 H).

#### Industrial Applicability


[0075] The novel functional PNA monomers in accordance with the present invention can be applied to the assembly of PNA that is used in, for example, gene therapy. Furthermore, in accordance with the present invention, a functional molecule can be efficiently introduced into PNA by two complementary synthetic routes B and C to the functional PNA monomers. The present invention can therefore be applied industrially, for example to an industrial synthesis of a functional PNA monomer unit using a Boc type monomer unit or a benzyloxycarbonyl- $\omega$ -amino acid derivative.

#### Claims


1. A compound represented by general formula (I) below:




(In the formula, A denotes



B denotes




R denotes H,  $\text{NO}_2$ ,  $\text{NH}_2$ ,  $\text{NHCbz}$ , Br, F, Cl or  $\text{SO}_3\text{Na}_2$ , and n is an integer of 1 to 4, wherein when A denotes





then B denotes




).

2. A process for producing a functional PNA monomer by reacting a t-butoxycarbonylaminoethylamine with a derivative of a functional molecule so as to incorporate the functional molecule into a PNA monomer, **characterised in that** the derivative of a functional molecule is an activated ester.
3. The production process according to Claim 2, **characterised in that** the activated ester has, on the carbonyl carbon forming the ester bond, a group represented by general formula (II) below.



(In the formula, A denotes




R denotes H, NO<sub>2</sub>, NH<sub>2</sub>, NHCbz, Br, F, Cl or SO<sub>3</sub>Na<sub>2</sub>, and n is an integer of 1 to 4.).

4. The production process according to either Claim 2 or Claim 3, **characterised in that** the activated ester has, on its carbonyl carbon, a pentafluorophenoxy group or a succinimidoxo group.
5. A process for producing the activated ester according to Claim 3, **characterised in that** it comprises reacting a carboxylic acid derivative of a functional molecule with a compound having a pentafluorophenoxy group or a succinimidoxo group.
6. A process for producing the carboxylic acid derivative of a functional molecule according to Claim 5, **characterised in that** it comprises reacting a derivative of the functional molecule with an aliphatic carboxylic acid.
7. A process for producing a functional PNA monomer from a functional molecule comprising:
  - producing a derivative of the functional molecule from the functional molecule,
  - producing a carboxylic acid derivative of the functional molecule from the derivative of the functional molecule,
  - producing an activated ester from the carboxylic acid derivative of the functional molecule, and

producing a functional PNA monomer from the activated ester, **characterised in that** it comprises one or more of a) to c) below:

- a) in the production of the carboxylic acid derivative of the functional molecule, reacting the derivative of the functional molecule with an aliphatic carboxylic acid;
- b) in the production of the activated ester, reacting the carboxylic acid derivative of the functional molecule with a compound having a pentafluorophenoxy group or a succinimidoxy group; and
- c) in the production of the functional PNA monomer, reacting a *t*-butoxycarbonylaminoethylamine with the activated ester derivative of the functional molecule.

8. A process for producing a functional PNA monomer by reacting a derivative of a functional molecule with an  $\omega$ -amino acid derivative represented by general formula (III) below, thereby incorporating the functional molecule into a PNA monomer, **characterised in that** the derivative of the functional molecule is an activated ester:




(In the formula,  $R^1$  denotes a hydrogen atom or a straight- or branched-chain  $C_1$  to  $C_5$  alkyl group, and  $m$  denotes an integer of 1 to 11.).

9. The production process according to Claim 8, **characterised in that** the activated ester has, on its carbonyl group forming an ester bond, a group represented by general formula (II) below, either directly or via an aliphatic chain or a peptide chain:



(In the formula, A denotes




R denotes H,  $\text{NO}_2$ ,  $\text{NH}_2$ ,  $\text{NHCbz}$ , Br, F, Cl or  $\text{SO}_3\text{Na}_2$ , and n is an integer of 1 to 4.).

10. The production process according to either Claim 8 or Claim 9, **characterised in that** the activated ester has, on its carbonyl carbon, a pentafluorophenoxy group or a succinimidoxy group.

11. A process for producing a functional PNA monomer from a functional molecule comprising producing an activated ester from the functional molecule and producing a functional PNA monomer from the activated ester, **characterised in that**


the production of the activated ester from the functional molecule comprises reacting *m*-methyl red with a compound that contains a succinimidoxy group, and/or

the production of the functional PNA monomer from the activated ester comprises reacting an  $\omega$ -amino acid derivative represented by general formula (III) below with a derivative of the functional molecule that is an activated ester, thereby incorporating the functional molecule into a PNA monomer:





10 (In the formula, R<sup>1</sup> denotes a hydrogen atom or a straight- or branched-chain C<sub>1</sub> to C<sub>5</sub> alkyl group, and m denotes an integer of 1 to 11.).

[Fig. 1]



[fig. 2]



## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/JP01/08120

## A. CLASSIFICATION OF SUBJECT MATTER

Int.Cl.<sup>7</sup> C07C245/08, 269/06, 271/20, C07D221/14, 311/08, 311/12, 311/80,  
475/14, 491/22, 495/04 // C09B69/10

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Int.Cl.<sup>7</sup> C07C, C07D, C07K, C09B, C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
CAPLUS (STN), REGISTRY (STN)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X Y	WO 98/37232 A2 (Georgia Tech Research Corporation), 27 August, 1998 (27.08.1998), Claims; pages 12 to 25; Figs. 8 to 11 & EP 968309 A2 & US 6117973 A & US 6225052 B1	1 3-6, 8-11
X Y	WO 92/20702 A1 (BUCHARDT, Ole et al.), 26 November, 1992 (26.11.1992), Claims; working examples 2 to 5; Fig. 6 & EP 586474 A1 & JP 6-506945 A & US 5714331 A & US 5736336 A & US 5766855 A & US 5786461 A & US 5977296 A & EP 1074559 A1 & US 6201103 B1	2, 7 3-6, 8-11

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

\* Special categories of cited documents:  
"A" document defining the general state of the art which is not  
considered to be of particular relevance  
"E" earlier document but published on or after the international filing  
date  
"L" document which may throw doubts on priority claim(s) or which is  
cited to establish the publication date of another citation or other  
special reason (as specified)  
"O" document referring to an oral disclosure, use, exhibition or other  
means  
"P" document published prior to the international filing date but later  
than the priority date claimed

"T" later document published after the international filing date or  
priority date and not in conflict with the application but cited to  
understand the principle or theory underlying the invention  
"X" document of particular relevance; the claimed invention cannot be  
considered novel or cannot be considered to involve an inventive  
step when the document is taken alone  
"Y" document of particular relevance; the claimed invention cannot be  
considered to involve an inventive step when the document is  
combined with one or more other such documents, such  
combination being obvious to a person skilled in the art  
"&" document member of the same patent family

Date of the actual completion of the international search  
28 November, 2001 (28.11.01)

Date of mailing of the international search report  
11 December, 2001 (11.12.01)

Name and mailing address of the ISA/  
Japanese Patent Office

Authorized officer

Facsimile No.

Telephone No.

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/JP01/08120

**Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

The special technical feature common to claims 1, 4-6, 10, and 11 (hereinafter referred to as "invention group (A)") is "a compound represented by the general formula (I)."

The special technical feature common to claims 2, 3, and 7-9 (hereinafter referred to as "invention group (B)") is "to react a functional molecule in the form of an active ester when the functional molecule is introduced into a PNA monomer."

There is hence no special technical feature common to invention group (A) and invention group (B). Therefore, these two invention groups are not considered to be so linked as to form a single general inventive concept.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest ☐ The additional search fees were accompanied by the applicant's protest.  
☒ No protest accompanied the payment of additional search fees.